

Two-photon decays of ρ and ω mesons and strong decays $\rho \rightarrow \eta\pi$ and $\omega \rightarrow \pi\pi$ in dense matter

A.E.Radzhabov and M.K.Volkov

*Bogoliubov Laboratory of Theoretical Physics,
Joint Institute for Nuclear Research, 141980 Dubna, Russia*

Abstract

Two-photon decays of ρ and ω mesons and strong decays $\rho \rightarrow \eta\pi$ and $\omega \rightarrow \pi\pi$ are described in the quark Nambu–Jona-Lasinio(NJL) model in dense medium. Decays of ρ -meson are especially interesting because they contain intermediate a_0 -meson which became a sharp resonance in the vicinity of phase transition of hadron matter into quark-gluon plasma. This leads to increasing ρ -meson decay widths and, as a result, two-photon decay of ρ -meson in medium became comparable with two-photon decays of scalar and pseudoscalar mesons in vacuum. Similar effect take place in the strong decay $\rho \rightarrow \eta\pi$.

Key words: scalar and vector mesons, strong and electromagnetic decays, nonzero chemical potential

PACS: 12.39.Ki, 13.20.Jf, 13.25.Jx

In the last years great attention is devoted to the study of the behavior of matter at extreme conditions. These conditions can be reached in the heavy-ion collision experiments which are performed now [1,2,3] and planned in future (SIS-300, LHC). Indeed, at high temperatures and/or densities which can appear in these experiments hadron matter can transit into quark-gluon plasma. The quark condensate and spontaneous breaking of chiral symmetry disappear and the chiral symmetry is restored. Let us emphasize that in this region many properties of elementary particles are noticeably changed. Especially it concerns the scalar mesons, their masses strongly decrease and approach the masses of the pseudoscalar mesons. The lifetime changes and strong decay channels can be closed. As a result, broad scalar resonances turn into sharp

Email addresses: aradzh@thsun1.jinr.ru (A.E.Radzhabov),
volkov@thsun1.jinr.ru (M.K.Volkov).

resonances. This can lead to the amplification of some processes mediated by the scalar resonances, such as $\pi\pi \rightarrow \gamma\gamma$ [4] and $\pi\pi \rightarrow \pi\pi$ [5].

Especially interesting is that in this domain new decay modes which are forbidden in vacuum can be opened up. It is two photon decays of vector mesons, strong decays $\rho \rightarrow \eta\pi$ and $\omega \rightarrow \pi\pi$, dilepton decays of scalar mesons and so on. This short paper is devoted to the description of the above mentioned vector meson decays.

For description of these processes we use the $U(3) \times U(3)$ NJL model with 'tHooft interaction [6,7,8,9,10]. Lagrangian of the model consists of two parts: $U(3) \times U(3)$ - symmetric four-quark interaction and 'tHooft determinant [11] which contains the six-quark interaction

$$\begin{aligned} \mathcal{L} = & \bar{q}(i\hat{\partial} - m^0)q + \frac{G_1}{2} \sum_{i=0}^8 \left[(\bar{q}\lambda_i q)^2 + (\bar{q}i\gamma_5\lambda_i q)^2 \right] - \frac{G_2}{2} \sum_{i=0}^8 (\bar{q}\gamma_\mu\lambda_i q)^2 - \\ & - K \{ \det[\bar{q}(1 + \gamma_5)q] + \det[\bar{q}(1 - \gamma_5)q] \}, \end{aligned} \quad (1)$$

where λ_i ($i=1,\dots,8$) are the Gell-Mann matrices and $\lambda^0 = \sqrt{\frac{2}{3}}\mathbf{1}$, with $\mathbf{1}$ being the unit matrix; $\bar{q} = \{\bar{u}, \bar{d}, \bar{s}\}$ are antiquark fields and q are quark fields; m^0 is a current quark mass matrix with diagonal elements m_u^0, m_d^0, m_s^0 ($m_u^0 \approx m_d^0$); G_1 and G_2 are the four-quark coupling constants in the scalar-pseudoscalar and vector channels; K is the six-quark coupling constant.

In this model a very important role is played by gap equations which describe spontaneous breaking of chiral symmetry, and connect current and constituent quark masses. These equations have the form

$$\begin{aligned} m_u &= m_u^0 + 8m_u G I_1^\Lambda(m_u) + 32m_u m_s K I_1^\Lambda(m_u) I_1^\Lambda(m_s), \\ m_s &= m_s^0 + 8m_s G I_1^\Lambda(m_s) + 32K \left(m_u I_1^\Lambda(m_u) \right)^2, \end{aligned} \quad (2)$$

where m_u and m_s are the constituent quark masses; $I_1^\Lambda(m)$ is the quadratically divergent integral describing the quark loop with one vertex, tadpole, with quark mass m and the cut-off parameter Λ .

It is possible to use different methods for investigation of the meson behavior in the hot and dense matter. The most popular one is the Matsubara technique [12]. However, for many applications it is more convenient to use an equivalent representation for the quark propagator derived in the "real time" formalism [13,14]




Fig. 1. Diagram describing the process $\rho \rightarrow \gamma\gamma$. Crossed diagram is not shown.

$$S(p, T, \mu) = (\hat{p} + m) \left[\frac{1}{p^2 - m^2 + i\epsilon} + i2\pi\delta(p^2 - m^2)(\theta(p^0)n(\mathbf{p}, \mu) + \theta(-p^0)n(\mathbf{p}, -\mu)) \right], \quad (3)$$

where

$$n(\mathbf{p}, \mu) = \left(1 + \exp \frac{E - \mu}{T} \right)^{-1} \quad (4)$$

is the Fermi-Dirac function for quarks, $E = \sqrt{\mathbf{p}^2 + m^2}$, T is the temperature and μ is the chemical potential. Here it is convenient to use the three-momentum cut-off Λ_3 . As a result, quadratically and logarithmically divergent integrals $I_1^{\Lambda_3}(m, T, \mu)$ and $I_2^{\Lambda_3}(m, T, \mu)$ take the form

$$I_1^{\Lambda_3}(m, T, \mu) = \frac{N_c}{(2\pi)^2} \int_0^{\Lambda_3} dp \frac{p^2}{E} (1 - n(\mathbf{p}, \mu) - n(\mathbf{p}, -\mu)),$$

$$I_2^{\Lambda_3}(m, T, \mu) = \frac{N_c}{2(2\pi)^2} \int_0^{\Lambda_3} dp \frac{p^2}{E^3} (1 - n(\mathbf{p}, \mu) - n(\mathbf{p}, -\mu)). \quad (5)$$

We assume that the model parameters G_1 , G_2 , K , m_u^0 , m_s^0 and Λ_3 do not depend on T and μ [14]. The dependence of constituent quark masses m_u , m_s on T and μ are calculated from the gap equations (2).

The model parameters in vacuum are fixed with the help of the following physical quantities: masses of π , K and ρ mesons, weak pion decay constant f_π , strong decay width of ρ -meson and mass difference of η - η' mesons. As a result, model parameters have the following values: $m_u = 280$ MeV, $m_u^0 = 2.1$ MeV, $m_s = 416$ MeV, $m_s^0 = 51$ MeV, $G_1 = 3.2$ GeV⁻², $G_2 = 16$ GeV⁻², $K = 4.6$ GeV⁻⁵, $\Lambda_3 = 1.03$ GeV.

The amplitude of the process $\rho \rightarrow \gamma\gamma$ has the form

$$A_{\rho \rightarrow \gamma\gamma}^{\alpha\mu\nu} = J_{\rho \rightarrow a_0}^\alpha D_{a_0} T_{a_0 \rightarrow \gamma\gamma}^{\mu\nu}, \quad (6)$$




Fig. 2. The behavior of the quark mass and meson masses M_π , M_{a_0} , $M_{a_0}^*$, M_ρ , M_η , M_σ as function of μ at $T = 20$ (left), 100 (right) MeV.

where $J_{\rho \rightarrow a_0}^\alpha$ describe the $\rho \rightarrow a_0$ transition, D_{a_0} is the a_0 -meson propagator, and $T_{a_0 \rightarrow \gamma\gamma}^{\mu\nu}$ is the amplitude of the a_0 two-photon decay (see fig.1).

Let us consider these elements in detail. The transition loop $\rho \rightarrow a_0$ has the form (see [15])

$$\begin{aligned}
 J_{\rho \rightarrow a_0}^\alpha &= C_{\rho \rightarrow a_0} \frac{1}{4} \int \frac{d^4 k}{(2\pi)^4} \text{tr}_s [S(k_-, T, \mu) \gamma^\alpha S(k_+, T, \mu)] = \\
 &C_{\rho \rightarrow a_0} 2m_u \int \frac{d^4 k}{(2\pi)^3} k^\alpha \frac{\delta(k_+^2 - m^2)}{k_-^2 - m^2} (n(\mathbf{k}_+, \mu) - n(\mathbf{k}_+, -\mu)) (\theta(k_+^0) - \theta(-k_+^0)), \\
 C_{\rho \rightarrow a_0} &= 4N_c g_{a_0} g_\rho, \quad k_\pm = k \pm \frac{p}{2}, \quad g_{a_0} = (4I_2^{\Lambda_3}(m, \mu, T))^{-1/2}, \quad g_\rho = \sqrt{6} g_{a_0},
 \end{aligned} \tag{7}$$

where g_{a_0} and g_ρ are the meson-quark coupling constants.

It is easy to see from eq. (7) that in vacuum, $\mu = 0$, this transition is forbidden. In medium this transition is nonzero and transversal over p , $p_\alpha J_{\rho \rightarrow a_0}^\alpha = 0$. Convolution of $J_{\rho \rightarrow a_0}^\alpha$ can be expressed in the form

$$|J|^2 = J_{\rho \rightarrow a_0}^\alpha J_{\alpha, \rho \rightarrow a_0} = -\frac{p^2}{\mathbf{p}^2} J_{\rho \rightarrow a_0}^0 J_{0, \rho \rightarrow a_0} \tag{8}$$

Only zero-component of $J_{\rho \rightarrow a_0}^\alpha$ is needed (see also [15])

$$\begin{aligned}
 J_{\rho \rightarrow a_0}^0 &= \frac{C_{\rho \rightarrow a_0} m}{(4\pi)^2 |\mathbf{p}|} \int_m^\infty dE \delta(E) [(2E + p^0) \ln(F_+) + (2E - p^0) \ln(F_-)] \\
 \delta(E) &= \frac{\sinh(\mu/T)}{\cosh(\mu/T) + \cosh(E/T)}, \quad F_\pm = \frac{p^2 \pm 2p_0 E + 2|\mathbf{p}|k}{p^2 \pm 2p_0 E - 2|\mathbf{p}|k}, \tag{9}
 \end{aligned}$$

where $k = \sqrt{E^2 - m^2}$.

The propagator of a_0 -meson has the form

$$D_{a_0} = \frac{1}{M_{a_0}^2 - M_\rho^2 - i\Gamma_{a_0}(M_\rho)M_{a_0}}$$

Note, ρ -meson mass weakly changes with T and μ [14](see also fig. 2). In the NJL model the mass of a_0 -meson is expressed as [16]

$$M_{a_0}^2 = g_{a_0}^2 \left[\frac{1}{G_{a_0}} - 8I_1^{\Lambda_3}(m_u) \right] + 4m_u^2$$

$$G_{a_0} = G_1 - 4Km_s I_1^{\Lambda_3}(m_s) \quad (10)$$

As a result, the a_0 -meson mass is lower than experimental

$$M_{a_0} \approx 800 \text{ MeV}, \quad M_{a_0}^{\text{exp}} = 984.7 \pm 1.2. \quad (11)$$

It is possible that a_0 -meson has an additional four-quark component [17,18,19]. In order to take it into account we introduce an additional term Δ in the mass formula for a_0 -meson

$$M_{a_0}^{*2} = M_{a_0}^2 + \Delta$$

The model part of the mass of a_0 -meson decreases with increasing T and μ and decreasing of order parameter m_u . It is natural to suppose that the additional term Δ also has a similar behavior. Particularly, we take Δ proportional to the square of the order parameter m_u , $\Delta \approx 4m_u^2$.

The decay width $a_0 \rightarrow \eta\pi$ is equal to

$$\Gamma_{a_0\eta\pi}(M_\rho) = \frac{g_{a_0\eta\pi}^2}{16\pi M_\rho} \sqrt{\left(1 - \left[\frac{M_\eta + M_\pi}{M_\rho}\right]^2\right) \left(1 - \left[\frac{M_\eta - M_\pi}{M_\rho}\right]^2\right)}$$

$$g_{a_0\eta\pi} = 2m_u g_{a_0} \sin \bar{\theta}, \quad \bar{\theta} = \theta - \theta_0, \quad (12)$$

where $\theta_0 \approx 35.3^\circ$ is the ideal mixing angle (ctg $\theta_0 = \sqrt{2}$) and θ is the singlet-octet mixing angle for pseudoscalar mesons.

The amplitude of the two-photon decay of a_0 meson has the form [20]

$$T_{a_0 \rightarrow \gamma\gamma}^{\mu\nu} = C_{a_0 \rightarrow \gamma\gamma} f_1(T, \mu) F^{\mu\nu}, \quad C_{a_0 \rightarrow \gamma\gamma} = \frac{2\alpha g_{a_0}}{3\pi m},$$

where $F^{\mu\nu}$ is the electromagnetic tensor, and function f_1 is [4]

Particle	Mass, MeV	$\Gamma_{\gamma\gamma}$, KeV
π^0	134.9766 ± 0.0006	$(7.8 \pm 0.5) \cdot 10^{-3}$
η	547.75 ± 0.12	1.29 ± 0.07
η'	957.78 ± 0.14	4.29 ± 0.15
σ	$400 - 1200$	a few
a_0	984.7 ± 1.2	0.3 ± 0.1
f_0	980 ± 10	$0.39^{+0.1}_{-0.13}$

Table 1

Two-photon decays of scalar and pseudoscalar mesons in vacuum [21].

$$f_1(T, \mu) = 1 - \frac{3}{2} m^2(T, \mu) \int_0^\infty dk \frac{k^3}{E^6(k)} \ln \left[\frac{E(k) + k}{E(k) - k} \right] [n(k; T, \mu) + n(k; T, -\mu)]$$

Now we can consider a full amplitude of the decay $\rho \rightarrow \gamma\gamma$. It is shown in figs.3, 4 for temperatures $T = 20, 100$ MeV as functions of $\mathbf{p}$, μ . It is easy to see that at some values of T and μ the two-photon decay width of ρ meson can be comparable with the two-photon decay width of scalar and pseudoscalar mesons in vacuum (see table 1).

Let us consider the process $\omega \rightarrow \gamma\gamma$. The diagrams for this process consist of $\omega \rightarrow \sigma$ transition, σ -meson propagator and two-photon decay of σ meson. The transition $\omega \rightarrow \sigma$ coincides with the transition $\rho \rightarrow a_0$. At large values of μ and T the σ -meson mass in NJL model can approximately be expressed in a very simple form

$$M_\sigma^2 \approx M_\pi^2 + 4m_u^2. \quad (13)$$

The width of σ is defined by the strong decay $\sigma \rightarrow \pi\pi$. The decay width can be expressed as

$$\Gamma_{\sigma\pi\pi}(M_\omega) = \frac{3}{2} \frac{g_{\sigma\pi\pi}^2}{16\pi M_\omega} \sqrt{1 - \frac{4M_\pi^2}{M_\omega^2}}, \quad g_{\sigma\pi\pi} = 2m_u g_{a_0} \cos \bar{\phi}, \quad \bar{\phi} = \theta_0 - \phi \quad (14)$$

where ϕ is the singlet-octet mixing angle for scalar mesons.

The amplitude of the two-photon decay of σ meson has the same form as $a_0 \rightarrow \gamma\gamma$. The difference is only in numerical factor, $C_{\sigma \rightarrow \gamma\gamma} = \frac{10\alpha g_{a_0}}{9\pi m}$.

The main difference with the decay $\rho \rightarrow \gamma\gamma$ is that the σ meson mass is always lower than the ω meson mass. In the process $\rho \rightarrow \gamma\gamma$ we deal with the sharp resonance. In the case $\omega \rightarrow \gamma\gamma$, with increasing T and μ the σ meson

mass will move away from the ω mass. Therefore, this σ meson propagator has no resonant behavior. As a result, this process is significantly smaller than $\rho \rightarrow \gamma\gamma$ as it shown in fig. 5.

Strong decays $\rho \rightarrow \pi\eta$ and $\omega \rightarrow \pi\pi$ ¹ can be considered in the same manner. It is necessary to consider the diagrams, which are similar to those in fig.1, where electromagnetic vertices are changed by strong ones. These decay widths² are given in fig.6 at temperature $T = 20$ MeV.

The main mechanism that opens decays in medium is the quark loop describing a transition of vector meson into scalar meson. In vacuum this loop is equal to zero. However, at finite values of the chemical potential and three-momentum this transition becomes possible [15].

In spite of this transition, intermediate scalar mesons σ and a_0 take part in two-photon vector meson decays. They play different roles in these processes. Indeed, a_0 -meson in the region near the critical μ values becomes a sharp resonance that leads to noticeable amplifications of decays $\rho \rightarrow \gamma\gamma$, $\rho \rightarrow \eta\pi$ whereas σ -meson taking part in the decays of ω -meson has other behavior. It has a non-resonance character and does not lead to amplification of processes. As a result the maximal value of the $\rho \rightarrow \gamma\gamma$ decay width at $T = 20$ MeV is equal to 2.8 KeV at $\mu = 337$ MeV and $|\mathbf{p}| = 360$ MeV. The maximal value of the $\omega \rightarrow \gamma\gamma$ decay width at $T = 20$ MeV is equal to 3.1 eV at $\mu = 330$ MeV and $|\mathbf{p}| = 410$ MeV.

Similar resonance effects with a_0 -meson are considered in the process $\eta\pi \rightarrow e^+e^-$ [23]. Note that σ -meson can also lead to resonance amplification of different processes where it participates as an intermediate state, such as $\pi\pi \rightarrow \gamma\gamma$ [4], $\pi\pi \rightarrow \pi\pi$ [5] and $\pi\pi \rightarrow e^+e^-$ [23].

In this paper, we do not consider nonresonance diagrams, such as a direct decay of ρ meson into two photon through the quark triangle diagram (this type of diagrams is considered in [24,25]). The diagrams with pion loops [26] also neglected because they have next order $1/N_c$ expansion. In future, we plan to consider non-resonance diagrams, process $\phi \rightarrow \gamma\gamma$ and dilepton processes with scalar mesons.

The authors should like to thank S.B. Gerasimov, Yu.L. Kalinovsky, O.V. Teryaev, and V.L. Yudichev for useful discussions. The work is partially supported by RFBR Grant no. 05-02-16699.

¹ The decay width $\omega \rightarrow \pi\pi$ in medium is calculated in nuclear matter in a hadronic model including mesons, nucleons and Δ isobars in [22].

² Note that the decay $\rho^0 \rightarrow \eta\pi^0$ is closed in vacuum whereas $\Gamma_{\omega \rightarrow \pi\pi} = 145 \pm 25$ KeV is only due to breaking of isotopic invariance [21].

References

- [1] C. Lourenco, Nucl. Phys. A698 (2002) 13, hep-ex/0105053.
- [2] C. Roland et al., Nucl. Phys. A698 (2002) 54, hep-ex/0105043.
- [3] M. Gazdzicki et al., J. Phys. G30 (2004) S701, nucl-ex/0403023.
- [4] M.K. Volkov et al., Phys. Lett. B424 (1998) 235, hep-ph/9706350.
- [5] D. Jido, T. Hatsuda and T. Kunihiro, Phys. Rev. D63 (2001) 011901, hep-ph/0008076.
- [6] M.K. Volkov, Fiz. Elem. Chast. Atom. Yadra 17 (1986) 433.
- [7] S. Klimt et al., Nucl. Phys. A516 (1990) 429.
- [8] S.P. Klevansky, Rev. Mod. Phys. 64 (1992) 649.
- [9] M.K. Volkov, Phys. Part. Nucl. 24 (1993) 35.
- [10] D. Ebert, H. Reinhardt and M.K. Volkov, Prog. Part. Nucl. Phys. 33 (1994) 1.
- [11] G. 't Hooft, Phys. Rev. Lett. 37 (1976) 8.
- [12] J.I. Kapusta, Finite Temperature Field Theory (Cambridge Univ. Press., 1999).
- [13] L. Dolan and R. Jackiw, Phys. Rev. D9 (1974) 3320.
- [14] D. Ebert et al., Int. J. Mod. Phys. A8 (1993) 1295.
- [15] D. Blaschke et al., Phys. Rev. C57 (1998) 438, nucl-th/9709058.
- [16] M.K. Volkov, M. Nagy and V.L. Yudichev, Nuovo Cim. A112 (1999) 225, hep-ph/9804347.
- [17] R.L. Jaffe, Phys. Rev. D15 (1977) 267.
- [18] N.N. Achasov, Nucl. Phys. A728 (2003) 425.
- [19] S.B. Gerasimov, Nucl. Phys. Proc. Suppl. 126 (2004) 210.
- [20] M.K. Volkov, Theor. Math. Phys. 101 (1994) 1473.
- [21] S. Eidelman et al., Phys. Lett. B592 (2004) 1.
- [22] W. Broniowski, W. Florkowski and B. Hiller, Eur. Phys. J. A7 (2000) 287, nucl-th/9905040.
- [23] O. Teodorescu, A.K. Dutt-Mazumder and C. Gale, Phys. Rev. C63 (2001) 034903, nucl-th/0008006.
- [24] O. Teryaev, Chin. J. Phys. 34 (1996) 1074.
- [25] V.V. Skalozub and A.Y. Tishchenko, hep-th/9907097.
- [26] H.A. Weldon, Phys. Lett. B274 (1992) 133.




Fig. 3. Two-photon decay width of ρ -meson with model mass of a_0 meson M_{a_0} (left) and with corrected mass of a_0 meson $M_{a_0}^*$ (right) as a function of μ and $|\mathbf{p}|$ for $T = 20$ MeV.




Fig. 4. Two-photon decay width of ρ -meson with model mass of a_0 meson M_{a_0} (left) and with corrected mass of a_0 meson $M_{a_0}^*$ (right) as a function of μ and $|\mathbf{p}|$ for $T = 100$ MeV.




Fig. 5. Two-photon decay width of ω -meson as a function of μ and $|\mathbf{p}|$ for $T = 20$ (left), 100 (right) MeV.




Fig. 6. Strong decay widths $\rho \rightarrow \pi\eta$ and $\omega \rightarrow \pi\pi$ at temperature $T = 20$ MeV.